

# Dental Trauma: PERMANENT TEETH

## BUMPED TOOTH

**What to do:**

- If slightly loose, eat a soft food diet and avoid habits such as thumbsucking.
- If tooth is loose, leave it alone and call us to get it treated.

**Managing Pain:**

- Give Tylenol or Motrin for pain

**When to call FPD:**

- Call our office to determine the need for an appointment

## CHIPPED TOOTH

**What to do:**

- LARGE CHIP: If you see pink in the middle or there is sensitivity, call our office. If you still have part of the tooth, keep it in milk until we can see you.
- SMALL CHIP: Avoid biting into the food with the tooth, eat a soft food diet, and avoid any habits such as thumbsucking.

**Managing Pain:**

- Give Tylenol or Motrin for pain
- Use cold compress or ice pack to reduce swelling

**When to call FPD:**

- Call our office to determine the need for an appointment

**Office: 317-598-9898**

**Emergency: 317-514-5157**

## INTRUDED/DISPLACED TOOTH (MOVED/PUSHED IN TOOTH)

**What to do:**

- Call our office to immediately

**Managing Pain:**

- Give Tylenol or Motrin for pain
- Use cold compress or ice pack to reduce swelling

**When to call FPD:**

- Call our office immediately



## AVULSED TOOTH (KNOCKED OUT TOOTH)

**What to do:**

- Do not touch the root. Rinse it off if it is dirty.
- Place tooth in milk or water. Never allow the tooth to dry.

**Managing Pain:**

- Give Tylenol or Motrin for pain

**When to call FPD:**

- Call our office immediately

## ABSCESS (INFECTION)/TOOTH PAIN

**What to do:**

- Give Tylenol or Motrin for pain
- Call our office to determine the need for an appointment

**Fishers Pediatric Dentistry**  
**Office: 317-598-9898**  
**Emergency: 317-514-5157**